

Determination of Iodine Value of an Oil

Practical 05 | Pharmaceutical Organic Chemistry II

Aim

To determine Iodine Value of an Oil, and interpret the recorded observations.

Principle

- The iodine value expresses halogen uptake as grams of iodine equivalent per 100 g of oil and is an index of unsaturation.
- A standardized titrant reacts with the analyte according to a defined stoichiometric relationship.
- The calculation must distinguish direct titration from a blank-corrected or back-titration procedure.

Materials and Apparatus

- Assigned oil, specified standardized reagents, blank reagents and oil-value glassware.
- Assigned test material, standardized titrant, specified solvent and indicator or endpoint detector.
- Burette, pipette, volumetric flask, conical flask and analytical balance.

Method

1. Identify the analyte form, the reaction and the endpoint specified for the exercise.
2. Document titrant concentration or factor and prepare a reagent blank where the method requires one.
3. Measure the prescribed test portion and dissolve or treat it by the assigned analytical method.
4. Titrate to the specified endpoint while recording initial and final burette readings.
5. Repeat the determination with independently measured portions and retain all readings.
6. Apply blank, dilution and stoichiometric corrections before reporting the result.

Observation Record

Complete this record using actual laboratory findings or the provided teaching data.

Observation / parameter	Record / finding
Oil mass	_____
Titrant concentration	_____
Blank volume	_____
Test volumes	_____
Calculated oil value	_____

Calculations and Interpretation

- Iodine value = $(B - S) \times N \times 12.69 / W$, in g iodine/100 g oil.
- Use the assigned iodometric method, thiosulfate normality and its prescribed blank; W is oil mass in g.

Worked Illustration

Illustration: $B - S = 4.00$ mL, $N = 0.100$ eq/L and $W = 0.500$ g give 10.15 g iodine/100 g oil.

The values in this illustration are for calculation practice.

Result

Record the actual finding, calculated value with units where relevant, and a conclusion supported by the observations.

Finding: _____

Conclusion: _____

Precautions

- Condition volumetric apparatus and avoid burette-tip air bubbles.
- Use the endpoint and solvent assigned to the analyte; do not transfer a factor from another salt or hydrate.
- Keep direct and back-titration signs consistent and report concordance.

Viva Questions

1. Why standardize a titrant?

Its actual concentration may differ from its nominal preparation concentration.

2. Why perform a reagent blank?

It measures reagent or solvent consumption unrelated to the analyte.

Continue Learning with FirstHope

Complete Notes on the FirstHope App

Complete subject and semesters are available on FirstHope app only.

- Get complete subject and semester notes in the FirstHope app.
- Available for B.Pharm, D.Pharm, M.Pharm and Pharm.D students.

[Click here to get the FirstHope app](#)

Join the FirstHope Telegram Channel

Join FirstHope Notes and keep up with channel updates.

[Click here to join Telegram](#)

Instagram

@firsthope.official

[Click here to follow Instagram](#)

YouTube Channel

FirstHope-Pharmacy / Live streams

[Click here to watch the YouTube channel](#)

WhatsApp Channel

Follow FirstHope on WhatsApp.

[Click here to join the WhatsApp channel](#)

FirstHope Website

firsthopenotes.com

[Click here to visit the website](#)