

Study of Stereotypy and Anti-Catatonic Activity

Practical 13 | Pharmacology I

Aim

To study Stereotypy and Anti-Catatonic Activity, and interpret the recorded observations.

Principle

- Repetitive-behaviour scores and posture-retention endpoints measure different behaviours; changes in general movement can confound both.
- This exercise uses a software/video simulation or an instructor-provided dataset.
- A response depends on the model, intervention, controls and observation period; comparisons must use the same endpoint.

Materials and Apparatus

- Assigned model/video and labelled completed pharmacology observations.
- Assigned pharmacology simulation/video and completed observation dataset.
- Computer, graphing or calculation software and a structured record sheet.

Method

1. Identify the simulated preparation, control group and response being measured.
2. Record the model settings and treatment labels supplied in the exercise.
3. Review the baseline or control trace before comparing the intervention traces.
4. Read the stated response endpoint using the same definition for every group.
5. Tabulate the completed simulation observations and calculate the assigned comparison.
6. Explain the mechanism, control response and any limitation of the model.

Observation Record

Complete this record using actual laboratory findings or the provided teaching data.

Observation / parameter	Record / finding
Treatment	_____
Stereotypy score	_____
Posture-retention time	_____
Motor observation	_____

Calculations and Interpretation

- Analyze each assigned behavioural endpoint separately and retain the scoring scale.

Result

Report the supplied simulated response, its defined comparison and the mechanism supported by the model. Label the observations as simulated.

Finding: _____

Conclusion: _____

Precautions

- Keep software units, treatment labels and endpoint definitions consistent.
- Use supplied simulation data and distinguish an illustration from an observed outcome.
- A model demonstrates an experimental principle; it does not establish an individual patient treatment.

Viva Questions

1. Why use a control response?

It separates the intervention effect from baseline variation in the simulated model.

2. Why define the endpoint before comparison?

Changing the endpoint after reviewing results can bias the interpretation.

Continue Learning with FirstHope

Complete Notes on the FirstHope App

Complete subject and semesters are available on FirstHope app only.

- Get complete subject and semester notes in the FirstHope app.
- Available for B.Pharm, D.Pharm, M.Pharm and Pharm.D students.

[Click here to get the FirstHope app](#)

Join the FirstHope Telegram Channel

Join FirstHope Notes and keep up with channel updates.

[Click here to join Telegram](#)

Instagram

@firsthope.official

[Click here to follow Instagram](#)

YouTube Channel

FirstHope-Pharmacy / Live streams

[Click here to watch the YouTube channel](#)

WhatsApp Channel

Follow FirstHope on WhatsApp.

[Click here to join the WhatsApp channel](#)

FirstHope Website

firsthopenotes.com

[Click here to visit the website](#)