

Limit Tests in Pharmaceutical Substances

Experiment 1 - Practical Notes

Name: _____ Roll no.: _____ Date: _____

For the heavy-metals component, use the explicitly assigned syllabus-era compendial method. A lead-specific limit test and a non-specific heavy-metals test are not equivalent; record the method and comparison basis. Modified chloride and sulphate procedures are included in their respective sections.

1. Chloride and Modified Chloride Limit Tests

Aim

To compare the impurity response of a test preparation with the prescribed standard under matched conditions.

Principle

Chloride reacts with silver ions to produce silver chloride turbidity: $\text{Ag}^+ + \text{Cl}^- \rightarrow \text{AgCl}$. The test and standard are compared after the same reagent additions and development conditions. The prescribed acidic medium helps reduce interference from ions such as carbonate.

Requirements

- Prescribed monograph for the test material, impurity standard and suitable matched comparison vessels.
- Silver nitrate test reagent, specified nitric acid and chloride-free vehicle.
- For the modified test: the material-specific approved pretreatment and recovery check.

Procedure

1. Read the monograph

Identify test mass, dissolution method, chloride-standard amount, reagent strengths and comparison time.

2. Prepare test and standard

Dissolve the test portion and prepare the standard with the stated amount of chloride. Use matched cylinders and the same final volume.

3. Develop turbidity

Add the prescribed acid and silver nitrate in the stated sequence. Mix equally and protect from inappropriate light exposure.

4. Compare

At the defined time compare turbidity under the prescribed viewing conditions; record less than, equal to or greater than the standard.

5. Perform the modified test

Use the prescribed substance-specific pretreatment, then carry out the matched comparison. Record how interference was removed and whether the impurity was retained.

Pharmaceutical Inorganic Chemistry

Modified Test: What Changes

- The precipitation principle remains the same; preparation is adapted to the named test material.
- For a carbonate-containing matrix, the prescribed acid treatment may remove carbonate with carbon dioxide evolution before the final comparison. This is a matrix example, not a universal modified recipe.
- For coloured or reactive material, use the monograph's approved decolourisation or other pretreatment and a suitable recovery/control check. Apply the same relevant treatment to the comparison preparation when specified.

Observations and Record

Item	Required record	Observation
Test portion / standard	Mass; impurity amount in standard	_____
Conditions	Reagents, final amount and development time	_____
Comparison	Test versus standard response	_____

Worked Example

The numbers below are illustrative; enter your own observations in the record.

- Illustrative limit calculation: a 2.0 g test portion has a stated teaching limit of 20 µg chloride/g. The comparator contains $2.0 \times 20 = 40$ µg chloride.
- A standard containing 20 µg/mL supplies 40 µg in 2.0 mL. These numbers demonstrate comparator preparation, not a universal IP chloride limit.

Result

The test response was ____ compared with the standard. Under valid matched conditions it ____ the prescribed limit; an invalid blank, standard or preparation requires repeat assessment.

Precautions

- Use chloride-free water and clean glassware; contamination can produce a false failure.
- Match reagent additions, final volume and reading time.
- Do not improvise pretreatment with an oxidiser or reducing agent; it may introduce chloride or alter recovery.
- Use the specific substance's prescribed limit; a teaching calculation does not set the permitted impurity concentration.

Viva Questions and Answers

1. What produces turbidity?

Silver chloride particles formed from chloride and silver ions.

2. Why use matched conditions?

The visual response depends on reagent concentration, time and optical conditions.

Pharmaceutical Inorganic Chemistry

3. Why can a modified test be necessary?

Colour, insolubility or chemical interference can obscure the ordinary comparison.

2. Sulphate and Modified Sulphate Limit Tests

Aim

To compare the impurity response of a test preparation with the prescribed standard under matched conditions.

Principle

Sulphate reacts with barium ions to form barium sulphate turbidity: $\text{Ba}^{2+} + \text{SO}_4^{2-} \rightarrow \text{BaSO}_4$. The prescribed acidic conditions and reagent system support a reproducible comparison. Carbonate and other matrix effects must not be allowed to produce unrelated turbidity.

Requirements

- Prescribed monograph for the test material, impurity standard and suitable matched comparison vessels.
- Specified barium chloride or barium-sulphate reagent system, acid and sulphate-free vehicle.
- Material-specific pretreatment for the modified test.

Procedure

1. Identify conditions

Record the material, test mass, sulphate-standard amount, reagent system and development time from the prescribed method.

2. Prepare the pair

Make test and standard to the same stated final amount in matched comparison vessels. Check that any insoluble material is handled as prescribed.

3. Develop turbidity

Add acid and the prescribed barium reagent in the specified order. Mix both equally.

4. Compare

At the fixed time compare the test turbidity with the standard under the same viewing conditions.

5. Apply the modification

Where the matrix interferes, use the approved material-specific preparation and verify that it has not removed or introduced sulphate.

Modified Test: Matrix Preparation

- A modified test keeps the sulphate-comparison objective but changes the preparation needed for a particular substance.
- Record the reason for modification: colour, poor dissolution or another specified interference.

Pharmaceutical Inorganic Chemistry

- If carbonate is present, the approved acid-treatment stage can remove it before barium addition. Strongly coloured materials need the prescribed preparation and controls; do not infer a method solely from the material's colour.

Observations and Record

Item	Required record	Observation
Test portion / standard	Mass; impurity amount in standard	_____
Conditions	Reagents, final amount and development time	_____
Comparison	Test versus standard response	_____

Worked Example

The numbers below are illustrative; enter your own observations in the record.

- Illustrative standard: 2.0 g material at a teaching limit of 50 µg/g requires 100 µg sulphate in the comparator.
- If the standard is 50 µg/mL, use 2.0 mL to supply 100 µg. Match the prescribed final volume and reagent additions.

Result

The test response was ____ compared with the standard. Under valid matched conditions it ____ the prescribed limit; an invalid blank, standard or preparation requires repeat assessment.

Precautions

- Do not replace the prescribed barium reagent system with an arbitrary solution strength.
- Keep volumes, mixing and time the same for test and standard.
- Use sulphate-free materials and avoid carryover from earlier tests.
- Use the specific substance's prescribed limit; a teaching calculation does not set the permitted impurity concentration.

Viva Questions and Answers

1. What is the insoluble product?

Barium sulphate.

2. Why may acidification be important?

It can suppress carbonate-related interference while providing the stated reaction conditions.

3. Does a darker starting colour mean more sulphate?

No; matrix colour is an interference, not a sulphate assay.

Pharmaceutical Inorganic Chemistry

3. Iron Limit Test

Aim

To compare the impurity response of a test preparation with the prescribed standard under matched conditions.

Principle

In the traditional teaching method, thioglycolic acid reduces iron as needed and forms a coloured iron complex in the prescribed alkaline medium. Citric acid assists the solution chemistry and limits precipitation. Colour is compared with a standard containing the allowed iron amount.

Requirements

- Prescribed monograph for the test material, impurity standard and suitable matched comparison vessels.
- Iron standard, thioglycolic-acid reagent, specified citric-acid and alkaline reagents.
- Iron-free vehicle and matched colour-comparison cylinders.

Procedure

1. Record the method

Identify the material, test portion, iron-standard amount and specified reagent strengths.

2. Prepare solutions

Dissolve or prepare the test as prescribed. Make the standard with the stated iron amount and a matched final volume.

3. Develop colour

Add the reagents in the specified order and adjust to the stated alkaline conditions. Mix and allow the same colour-development time for both.

4. Inspect controls

Check that reagent blanks and the standard behave acceptably. A precipitate or obscuring starting colour may invalidate comparison.

5. Compare

Compare the developed colour, commonly purple in this teaching method, and record the relation to the standard.

Observations and Record

Item	Required record	Observation
Test portion / standard	Mass; impurity amount in standard	_____
Conditions	Reagents, final amount and development time	_____
Comparison	Test versus standard response	_____

Pharmaceutical Inorganic Chemistry

Worked Example

The numbers below are illustrative; enter your own observations in the record.

- Illustrative standard amount: 1.0 g test portion at 10 µg iron/g requires 10 µg iron.
- A 5 µg/mL standard supplies this amount in 2.0 mL. A matched colour comparison is still required; arithmetic alone is not a test result.

Result

The test response was ____ compared with the standard. Under valid matched conditions it ____ the prescribed limit; an invalid blank, standard or preparation requires repeat assessment.

Precautions

- Use iron-free reagents and avoid metal contamination.
- The alkaline condition must match the method; uncontrolled pH changes can affect colour or precipitation.
- Do not use the colour response as a quantitative iron assay unless the method has been validated for that purpose.
- Use the specific substance's prescribed limit; a teaching calculation does not set the permitted impurity concentration.

Viva Questions and Answers

1. What is the role of thioglycolic acid?

It supports reduction and formation of the coloured iron complex.

2. Why is citric acid included?

It assists complexation and helps prevent unwanted precipitation under the method conditions.

3. What does a limit test report?

Whether the impurity response is within the prescribed comparator limit, under valid conditions.

4. Heavy Metals - Assigned Compendial Limit Test

Aim

To compare the supplied heavy-metals limit-test response with its prescribed standard.

Principle

The syllabus-era non-specific heavy-metals test assesses a method-defined response, commonly a coloured sulphide comparison under specified conditions. It is not identical to a lead-specific test or to a modern element-specific instrumental procedure.

Requirements

- Instructor-provided test and standard comparison records or the prescribed supervised compendial demonstration.

Pharmaceutical Inorganic Chemistry

Procedure

1. Identify the method

Record monograph, edition, test preparation and comparison standard basis.

2. Check suitability

Identify the assigned digestion/preparation and matrix-control requirements; use the approved demonstration rather than improvising reagent handling.

3. Compare responses

Read the test and standard under the same specified visual conditions and record actual colour/intensity.

4. Conclude

State whether the valid method-defined test response exceeds the assigned standard; do not name or quantify an individual metal from a non-specific response.

Observations and Record

Comparison	Observation
Assigned method / material	_____
Standard response	_____
Test response	_____
Control / limit conclusion	_____

Result

Record the completed observations, calculations and conclusion with units. State any limitation of the observation.

Precautions

- Use only the approved method and facilities for hazardous reagents.
- Do not substitute a lead-specific test for a non-specific heavy-metals test.

Viva Questions and Answers

1. What does the comparison establish?

Conformity with the assigned method-defined response limit.

2. Does it identify each metal?

No. A non-specific visual test does not provide element-specific quantities.

5. Arsenic Limit Test

Aim

To compare the impurity response of a test preparation with the prescribed standard under matched conditions.

Pharmaceutical Inorganic Chemistry

Principle

The classical Gutzeit principle detects arsenic through a stain produced in a controlled gas-generation apparatus. The generated arsenic-containing gas is highly toxic, so this practical is taught through a sealed authorised demonstration, supplied records or a simulation. No gas-generation recipe is part of these notes.

Requirements

- Prescribed monograph for the test material, impurity standard and suitable matched comparison vessels.
- Instructor-operated closed demonstration or approved video and comparator stain record.
- Method-specific standard information and observation sheet.

Procedure

1. Identify the principle

Label the closed reaction section, interference-control section and stain-detection section on the supplied apparatus diagram.

2. Check demonstration records

Record test-portion identity, standard arsenic amount, controlled exposure period and the instructor's validity checks.

3. Observe without intervention

Study the sealed demonstration or video. Students do not prepare gas-generating reagents, open the apparatus or handle detector material.

4. Compare stains

Use the supplied valid test and standard stain records. Compare colour intensity under the prescribed viewing conditions.

5. Report

State the relation to the standard and distinguish a classical teaching demonstration from a validated modern instrumental arsenic assay.

Observations and Record

Item	Required record	Observation
Test portion / standard	Mass; impurity amount in standard	_____
Conditions	Reagents, final amount and development time	_____
Comparison	Test versus standard response	_____

Worked Example

The numbers below are illustrative; enter your own observations in the record.

Pharmaceutical Inorganic Chemistry

- Illustrative comparator arithmetic only: 1.0 g material at a stated 2 µg/g limit requires 2 µg arsenic in the comparison.
- If a supplied standard is 1 µg/mL, 2.0 mL contains 2 µg. This is a paper calculation, not an instruction to generate gas.

Result

The test response was ____ compared with the standard. Under valid matched conditions it ____ the prescribed limit; an invalid blank, standard or preparation requires repeat assessment.

Precautions

- Use the authorised closed demonstration or simulation only.
- Do not open, assemble or run a gas-generation experiment from a textbook description.
- Hazardous residues and detector materials are handled by authorised staff.
- Use the specific substance's prescribed limit; a teaching calculation does not set the permitted impurity concentration.

Viva Questions and Answers

1. What is compared?

The developed test stain against the prescribed standard stain.

2. Why are matched conditions needed?

Stain intensity depends on the controlled exposure and apparatus conditions.

3. Does this test identify every arsenic species separately?

No; the classical limit response is not a complete species-specific analysis.

Continue Learning with FirstHope

Complete Notes on the FirstHope App

Complete subject and semesters are available on FirstHope app only.

- Get complete subject and semester notes in the FirstHope app.
- Available for B.Pharm, D.Pharm, M.Pharm and Pharm.D students.

[Click here to get the FirstHope app](#)

Join the FirstHope Telegram Channel

Join FirstHope Notes and keep up with channel updates.

[Click here to join Telegram](#)

Instagram

@firsthope.official

[Click here to follow Instagram](#)

YouTube Channel

FirstHope-Pharmacy / Live streams

[Click here to watch the YouTube channel](#)

WhatsApp Channel

Follow FirstHope on WhatsApp.

[Click here to join the WhatsApp channel](#)

FirstHope Website

firsthopenotes.com

[Click here to visit the website](#)