

Complexation and protein binding

Introduction to Complexation and Protein Binding

Complexation

- Complexation is a chemical process where molecules or ions combine to form a stable complex.
- It typically involves:
 - A. Central Metal Ion:** Usually a metal (e.g., Fe, Cu, Zn) that acts as an electron acceptor.
 - B. Ligands:** Molecules or ions that donate electron pairs to the metal, forming coordinate covalent bonds. Ligands can be small ions (like Cl^-) or larger molecules (e.g., ethylenediamine).
- Complexation is key in coordination chemistry, influencing the structure and stability of complexes, which can vary from linear to complex geometries (e.g., octahedral).
- It plays a significant role in processes like catalysis and metal ion transport in biological systems.


Protein Binding

- Protein binding involves the interaction of drugs or molecules with proteins in biological systems, particularly in the bloodstream.
 - Plasma Proteins:** Proteins like albumin bind to various molecules, affecting their availability.
 - Drug Impact:** When bound to proteins, drugs are often inactive; only the unbound fraction is pharmacologically active.
 - Pharmacokinetics/Pharmacodynamics:** Protein binding influences a drug's movement, duration, and elimination in the body, impacting its therapeutic effects.

Classification of Complexation

- The classification of complexation, as shown in the image, is structured into three main categories.

Here's a detailed explanation:



1. Metal Ion Complexes/Coordinate Complexes

- These are complexes formed by metal ions binding with various ligands.
- This category includes:
 - **Inorganic Type:** These complexes involve metal ions forming bonds with inorganic ligands.
 - **Chelates:** A specific type of complex where metal ions bind with a ligand that has multiple binding sites, creating a ring structure. Chelates are stable due to the formation of multiple bonds with the metal ion.
 - **Olefin Type:** These complexes involve metal ions binding with olefins (unsaturated hydrocarbons containing a carbon-carbon double bond).
 - **Aromatic Type:** Complexes formed when metal ions bind with aromatic compounds, which are compounds containing a conjugated pi-electron system, like benzene.

2. Organic Molecular Complexes

- These complexes involve organic molecules interacting with other compounds.
- They are further classified into:
 - **Quinhydrone Type:** Complexes formed by quinhydrone, which is a compound consisting of a 1:1 mixture of quinone and hydroquinone.
 - **Picric Acid Type:** Complexes formed with picric acid, which is a nitroaromatic compound that acts as a ligand.

- **Caffeine and Other Drug Complexes:** These complexes involve drugs like caffeine interacting with other molecules, often forming complexes through hydrogen bonding or other interactions.
- **Polymer Type:** Complexes involving polymers, which are large molecules composed of repeated subunits, forming bonds with smaller molecules or ions.

3. Inclusion/Exclusion Compounds

- These are complexes where one molecule (the host) forms a cavity or structure that encapsulates another molecule (the guest).
- This category includes:
 - **Clathrate:** A compound in which molecules are physically trapped within the lattice structure of another compound without any chemical bonding between them.
 - **Channel Lattice Type:** Complexes where the host molecule creates a channel-like structure, allowing the guest molecule to be included.
 - **Layer Type:** These involve layers of host molecules trapping the guest molecules between them.
 - **Monomolecular Type:** Single host molecules encapsulating guest molecules.
 - **Macromolecular Type:** Large host molecules, often macromolecules like cyclodextrins, that include smaller guest molecules within their structure.

Types of Ligands

1. **Monodentate Ligands:** Bind to the central atom through a single donor atom.
2. **Bidentate Ligands:** Have two donor atoms that can bind to the central atom simultaneously.
3. **Polydentate Ligands:** Have three or more donor atoms that can attach to the central atom.
4. **Chelating Ligands:** Form one or more rings by binding to the central atom through multiple donor atoms.
5. **Ambidentate Ligands:** Have two or more donor atoms, but only one binds to the central atom at a time.
6. **Bridging Ligands:** Simultaneously bind to two or more central atoms, forming a bridge between them.

Applications of Complexation

Complexation plays a crucial role in various scientific and industrial applications:

1. Pharmaceuticals

- **Drug Delivery:** Complexes can enhance the solubility, stability, and bioavailability of drugs.
- **Metal-based Drugs:** Complexes like cisplatin are used in cancer therapy.



FREE PREVIEW

[Click here to visit our website](#)

PHARMACY PREMIUM NOTES

Complete subject
and semesters are
available on
FirstHope app only.



[Scan or click here](#)

Click here to get the app



Available on Google Play

JOIN THE FIRSTHOPE COMMUNITY

Join our Telegram channel

Click here to join Telegram



[Scan or click here](#)

@FirstHopeNotes

Instagram

@firsthope.official

[Click here to follow >](#)

YouTube channel

FirstHope-Pharmacy / Live streams

[Click here to watch >](#)

WhatsApp channel

Follow FirstHope on WhatsApp

[Click here to join >](#)

FirstHope website

firsthopenotes.com

[Click here to visit >](#)

B.Pharm / D.Pharm / M.Pharm / Pharm.D

Preview notes. Complete subjects and semesters in the FirstHope app.