

Pharmacokinetics

Pharmacokinetics: Definition and Introduction

Definition

- Pharmacokinetics (PK) is the branch of pharmacology that deals with the quantitative study of how a drug moves into, through, and out of the body over time.
- It answers the question: **"What does the body do to the drug?"**
- Pharmacokinetics focuses on four primary processes, collectively referred to as **ADME**:
 - **Absorption**
 - **Distribution**
 - **Metabolism**
 - **Excretion**

Importance of Pharmacokinetics

- Helps determine proper dosage regimens, frequency of administration, and the optimal route of drug delivery.
- Aids in predicting drug-drug interactions, avoiding toxicity, and optimizing therapeutic efficacy.

Key Aspects of Pharmacokinetics (ADME)

1. Absorption

- The process of a drug entering the **bloodstream** from its site of administration.

2. Distribution

- The dispersion of the drug throughout the **fluids and tissues** of the body.

3. Metabolism (Biotransformation)

- The **biochemical modification** of the drug, mainly occurring in the **liver**.

4. Excretion

- The removal of the drug and its metabolites, primarily via the **kidneys (urine)** or **bile (feces)**.

Mathematical Model in Pharmacokinetics

Overview

- A mathematical model in pharmacokinetics is a **quantitative framework** that describes how a drug is absorbed, distributed, metabolized, and eliminated (ADME) in the body over time.

- These models use equations and variables to simulate drug concentrations at various sites (like blood or tissues) to predict outcomes and guide dosage regimens.

Why Use Mathematical Models?

- To predict drug concentrations at any time point.
- To simulate different dosing scenarios (single dose vs. multiple doses).
- To optimize therapeutic efficacy and minimize toxicity.
- To aid in designing clinical trials and interpreting pharmacokinetic data.

Types of Mathematical Models

There are two broad categories:

A. Compartmental Models

- These treat the body as one or more "compartments" where the drug moves between.
- Simple but useful for most clinical applications.

B. Physiologically Based Pharmacokinetic (PBPK) Models

- These are complex models that incorporate real physiological parameters like organ sizes, blood flow rates, etc.
- Used in regulatory submissions and detailed simulations.

Basic Mathematical Elements

- **Differential Equations:** These describe the rate of change of drug concentration over time.
- **Rate Constants (k):** Represent processes such as absorption (k_a), elimination (k_e), and distribution (k_{12} , k_{21}).
- **Volume of Distribution (Vd):** The theoretical volume in which the drug would need to be uniformly distributed to produce the observed concentration.
- **Clearance (Cl):** The rate at which a drug is removed from the body.

Example Equation

- For a single-compartment model with first-order elimination:

$$C(t) = C_0 \cdot e^{-k_e t}$$

- Where:
 - $C(t)$ = drug concentration at time t
 - C_0 = initial concentration
 - k_e = elimination rate constant

Reaction Kinetics: Rate, Rate Constants, and Order

Basic Concepts

A. Rate of Reaction:

- Measures how fast reactants are converted into products. Influenced by **temperature**, **concentration**, **catalysts**, and **surface area**.

B. Rate Constant (k):


- A constant specific to a given reaction at a certain temperature.

C. Order of Reaction:

- Indicates how the **reaction rate depends on the concentration** of reactants.

General Rate Equation:

- For a reaction:



$$\text{Rate} = k[A]^m[B]^n$$

- **m & n:** Orders of the reaction with respect to A and B, respectively.
- **Overall Reaction Order** = m + n

Types of Reaction Kinetics

A. Zero-Order Kinetics

- Rate is **independent of reactant concentration**.
- Drug concentration **decreases linearly over time**.

Rate Equation:

$$\text{Rate} = k$$

Half-life ($t_{1/2}$):

$$t_{1/2} = \frac{[A]_0}{2k}$$

- Half-life **depends on initial concentration**.

B. First-Order Kinetics

- Rate is **directly proportional** to the concentration of the drug.
- Drug elimination follows an **exponential decline**.

Rate Equation:

$$\text{Rate} = k[A]$$

Half-life ($t_{1/2}$):

$$t_{1/2} = \frac{0.693}{k}$$

- Half-life is **constant**, regardless of concentration.

Plasma Drug Concentration-Time Profile / Drug Levels in Blood

- Plasma drug concentration-time profiles (often referred to as **drug levels in blood**) represent how the concentration of a drug in the bloodstream changes over time.
- This data is essential for evaluating the **ADME** phases (Absorption, Distribution, Metabolism, and Elimination) and guiding clinical decisions regarding **dosage** and **therapeutic drug monitoring (TDM)**.

Phases of Drug in Blood

1. Absorption Phase


- Occurs after administration (especially non-IV routes).
- Drug levels increase as the drug enters systemic circulation.

2. Distribution Phase

- Drug begins to move from the blood into various tissues.
- Plasma levels may drop even though the total body drug amount is still high.

3. Elimination Phase

- Metabolism and excretion lower the drug concentration over time.



- Represents **total drug exposure** over time.
- Helps in determining **half-life ($t_{1/2}$)**, **clearance (Cl)**, and **volume of distribution (Vd)**.

Key Pharmacokinetic Parameters

- **C_{max}**: Peak plasma concentration achieved after administration.
- **T_{max}**: Time taken to reach C_{max}.
- **AUC (Area Under the Curve)**: Total drug exposure over time.
- **Half-life ($t_{1/2}$)**: Time required for plasma concentration to reduce by 50%.
- **Clearance (Cl)**: Volume of plasma cleared of the drug per unit time.
- **Volume of Distribution (Vd)**: Theoretical volume linking the plasma concentration to the total amount of drug in the body.

Importance in Clinical Settings

- **Therapeutic Drug Monitoring (TDM)**: Especially for drugs with narrow therapeutic windows (e.g., digoxin, lithium).
- **Dose Adjustment**: Based on individual metabolism, age, disease state, or drug interactions.
- **Bioavailability Assessment**: Compares drug levels between different formulations.

Drug Monitoring Techniques

- **Blood Sampling**: At specific time intervals.
- **Chromatographic Methods**: HPLC, LC-MS used for measuring drug levels.
- **Non-invasive Alternatives**: Under development (e.g., saliva, sweat analysis).

Pharmacokinetic Model

Definition

- A **pharmacokinetic model** is a conceptual and mathematical tool used to describe and predict the movement of drugs within the body over time.
- These models provide a simplified representation of complex physiological processes, aiming to quantify **absorption**, **distribution**, **metabolism**, and **excretion** (ADME) using parameters derived from experimental data.

Purpose of Pharmacokinetic Models

- To simulate how a drug behaves in the human body.
- To predict plasma and tissue concentrations over time.
- To estimate important pharmacokinetic parameters like **half-life**, **volume of distribution**, and **clearance**.



FREE PREVIEW

[Click here to visit our website](#)

PHARMACY PREMIUM NOTES

Complete subject
and semesters are
available on
FirstHope app only.



[Scan or click here](#)

Click here to get the app



Available on Google Play

JOIN THE FIRSTHOPE COMMUNITY

Join our Telegram channel

Click here to join Telegram



[Scan or click here](#)

@FirstHopeNotes

Instagram

@firsthope.official

[Click here to follow >](#)

YouTube channel

FirstHope-Pharmacy / Live streams

[Click here to watch >](#)

WhatsApp channel

Follow FirstHope on WhatsApp

[Click here to join >](#)

FirstHope website

firsthopenotes.com

[Click here to visit >](#)

B.Pharm / D.Pharm / M.Pharm / Pharm.D

Preview notes. Complete subjects and semesters in the FirstHope app.