

Pain management

Pain Management

Overview:

- Pain management refers to the strategies and interventions used to alleviate or control pain.
- It encompasses a wide range of pharmacologic and non-pharmacologic methods to treat acute, chronic, neuropathic, or nociceptive pain.
- In pharmacotherapeutics, the focus is on understanding the types of pain, underlying mechanisms (pathophysiology), and the pharmacological agents used for pain relief, their mechanisms of action, indications, side effects, and interactions.


Types of Pain (from a therapeutic perspective):

1. **Nociceptive pain** – Arises from tissue injury or inflammation. Subtypes include:
 - **Somatic pain:** Originates from skin, muscles, or bones. Well-localized.
 - **Visceral pain:** Originates from internal organs. Poorly localized, often described as deep, squeezing, or cramping.
2. **Neuropathic pain** – Arises from damage to or dysfunction of the nervous system.
 - Examples: diabetic neuropathy, postherpetic neuralgia, phantom limb pain.
3. **Mixed pain** – Has both nociceptive and neuropathic components (e.g., cancer pain).

Goals of Pharmacologic Pain Management:

- Relieve pain to improve quality of life.
- Minimize drug side effects.
- Restore function.
- Prevent pain recurrence (especially in chronic pain).
- Avoid or manage tolerance, dependence, or addiction (especially with opioids).

Analgesic Drug Classes:



1. Non-opioid Analgesics:

A. Acetaminophen (Paracetamol):

- **Mechanism:** Inhibits prostaglandin synthesis in the CNS.
- **Uses:** Mild to moderate pain, fever.
- **Limitations:** Hepatotoxicity at high doses.

B. NSAIDs (e.g., Ibuprofen, Naproxen, Diclofenac):

- **Mechanism:** Inhibit COX-1 and/or COX-2 → decreased prostaglandin production.
- **Uses:** Inflammatory pain, arthritis, musculoskeletal pain.
- **Adverse effects:** Gastric ulcers, renal impairment, cardiovascular risks.

2. Opioid Analgesics:

- **Examples:** Morphine, Fentanyl, Oxycodone, Tramadol.
- **Mechanism:** Bind to opioid receptors (μ , κ , δ) in CNS and periphery to inhibit pain transmission.
- **Uses:** Moderate to severe pain (post-surgical, cancer, palliative care).
- **Adverse effects:** Sedation, constipation, respiratory depression, tolerance, dependence, addiction.
- **Monitoring required** for dosage adjustment, dependence risk, especially in chronic use.

3. Adjuvant Analgesics:

- **Antidepressants (e.g., amitriptyline, duloxetine):** Used in neuropathic pain.
- **Anticonvulsants (e.g., gabapentin, pregabalin):** Effective for nerve pain.

- **Corticosteroids:** Useful in inflammatory or cancer-related pain.
- **Muscle relaxants:** Used in musculoskeletal pain with muscle spasms.

4. Topical Agents:

- **Lidocaine patches, capsaicin cream:** Used for localized neuropathic pain.

5. Emerging Therapies:

- NMDA receptor antagonists (e.g., ketamine).
- Cannabinoids (under evaluation in some countries).
- Monoclonal antibodies (e.g., anti-NGF).

Approaches to Pain Management (Pharmacotherapeutic Strategy):

1. WHO Analgesic Ladder (originally for cancer pain):

- **Step 1:** Non-opioid ± adjuvant
- **Step 2:** Weak opioid ± non-opioid ± adjuvant
- **Step 3:** Strong opioid ± non-opioid ± adjuvant

2. Individualized Treatment Plan:

- Assess pain severity (e.g., using visual analog scale).
- Consider patient-specific factors (age, comorbidities, renal/hepatic function, risk of abuse).
- Regular reassessment and adjustment.

3. Multimodal Analgesia:

- Using different classes of drugs targeting multiple pain pathways to enhance pain relief and minimize side effects.

Pain Pathways

Overview:

- Understanding pain pathways is essential in pharmacotherapeutics, as most analgesic drugs act on one or more components of these pathways.
- Pain perception (nociception) involves a series of complex neural events and signaling mechanisms.

Phases of Nociception:

1. Transduction:

- Nociceptors (pain receptors) are activated by noxious stimuli (thermal, mechanical, chemical).

- Chemical mediators involved include prostaglandins, bradykinin, histamine, serotonin, and substance P.
- These lower the threshold of nociceptors, making them more sensitive to stimuli.

2. Transmission:

- Pain signals are transmitted from the periphery to the CNS through afferent nerve fibers:
 - **A-delta fibers:** Fast, myelinated, transmit sharp pain.
 - **C fibers:** Slow, unmyelinated, transmit dull, aching pain.
- Signals travel to the **dorsal horn** of the spinal cord, then to the **thalamus**, and finally to the **cerebral cortex**.

3. Perception:

- The conscious awareness and interpretation of pain occur in the brain (somatosensory cortex, limbic system, frontal cortex).
- Emotional and psychological responses are integrated here.

4. Modulation:

- The CNS can enhance or inhibit pain transmission.
- **Descending inhibitory pathways** from the brainstem (periaqueductal gray, raphe nuclei) release neurotransmitters (e.g., serotonin, norepinephrine, endogenous opioids) that inhibit pain signal transmission in the spinal cord.
- Pharmacologically, this is where antidepressants and opioids can act.

Key Neurotransmitters in Pain Pathways:

1. Excitatory:

- Glutamate
- Substance P
- Bradykinin
- Prostaglandins

2. Inhibitory:

- GABA
- Endorphins, enkephalins (endogenous opioids)
- Serotonin
- Norepinephrine

Sites of Drug Action in Pain Pathways:

Site	Drug Classes Acting Here
------	--------------------------

Peripheral nociceptors	NSAIDs (block prostaglandins)
Spinal cord dorsal horn	Opioids (inhibit neurotransmitter release), local anesthetics
Descending inhibitory pathways	Antidepressants (increase serotonin/norepinephrine)
Cortical perception	Sedatives, anxiolytics, cognitive-behavioral therapy (not pharmacological but influences perception)

Pathophysiological Considerations:

- **Peripheral sensitization:** Increased responsiveness of nociceptors due to inflammatory mediators.
- **Central sensitization:** Increased excitability of spinal cord neurons → chronic pain states like fibromyalgia.
- Pharmacotherapeutic agents aim to target these mechanisms (e.g., NMDA receptor antagonists like ketamine can reduce central sensitization).

Neuralgias

Overview:

- **Neuralgia** refers to pain that occurs along the course of a nerve. It is a form of **neuropathic pain**, which results from damage or dysfunction in the peripheral or central nervous system.
- In pharmacotherapeutics, neuralgias present a particular challenge as they often do not respond well to conventional analgesics like NSAIDs or acetaminophen.

Common Types of Neuralgias:

1. Trigeminal Neuralgia (Tic Douloureux):

- Severe, stabbing, shock-like pain along the **trigeminal nerve (CN V)**, typically affecting one side of the face.
- Triggers: Touch, chewing, talking, brushing teeth.

2. Postherpetic Neuralgia:

- Follows **herpes zoster (shingles)** infection.
- Persistent burning or stabbing pain in the affected dermatome (often thoracic).

3. Occipital Neuralgia:

- Sharp pain in the upper neck, back of the head, or behind the eyes due to irritation or injury of the occipital nerves.

4. Glossopharyngeal Neuralgia:

- Pain in the throat, tonsils, tongue, or ear related to the glossopharyngeal nerve (CN IX).



FREE PREVIEW

[Click here to visit our website](#)

PHARMACY PREMIUM NOTES

Complete subject
and semesters are
available on
FirstHope app only.



[Scan or click here](#)

Click here to get the app



Available on Google Play

JOIN THE FIRSTHOPE COMMUNITY

Join our Telegram channel

Click here to join Telegram



[Scan or click here](#)

@FirstHopeNotes

Instagram

@firsthope.official

[Click here to follow >](#)

YouTube channel

FirstHope-Pharmacy / Live streams

[Click here to watch >](#)

WhatsApp channel

Follow FirstHope on WhatsApp

[Click here to join >](#)

FirstHope website

firsthopenotes.com

[Click here to visit >](#)

B.Pharm / D.Pharm / M.Pharm / Pharm.D

Preview notes. Complete subjects and semesters in the FirstHope app.